

Jun Ren¹ and James R. Sowers²

Application of a Novel Curcumin Analog in the Management of Diabetic Cardiomyopathy

Diabetes 2014;63:3166–3168 | DOI: 10.2337/db14-0863

Diabetes mellitus is becoming an epidemic health threat and represents one of the most prevalent chronic non-communicable disorders. Cardiovascular complications are considered the leading cause of death for diabetic patients. Diabetes leads to undesired changes in cardiac structure and function, a condition commonly known as diabetic cardiomyopathy, which occurs independent of macro- and microvascular comorbidities in diabetes (1,2). Both systolic and diastolic dysfunctions have been demonstrated in diabetic hearts including prolonged duration of contraction and relaxation, reduced velocity of contraction and relaxation, and depressed myocardial contractility (1,3). In particular, clinical evaluation using electrocardiogram and echocardiography has revealed substantial functional changes in diabetic hearts, including shorter left ventricular ejection time, prolonged pre-ejection duration, increased wall stiffness, decreased fractional shortening, decreased rate of left ventricular filling, and increased action potential duration (4,5). To date, a plethora of cellular and molecular mechanisms have been postulated for the onset and development of diabetic cardiomyopathy, including reduced energy production due to decreases in mitochondrial respiration and pyruvate dehydrogenase activity, accumulation of reactive oxygen species, oxidative stress, apoptosis, impaired autophagy, and malfunction of cardiac contractile and intracellular Ca^{2+} regulatory proteins such as myosin, sarco/endoplasmic reticulum Ca^{2+} -ATPase (SERCA), and $\text{Na}^+/\text{Ca}^{2+}$ exchanger (1–3,5,6). The high morbidity and mortality for diabetic cardiomyopathy warrant aggressive clinical management involving lifestyle modification, control of glucose and lipid abnormalities, and treatment of hypertension and coronary artery diseases, if present. The commonly used therapeutic regimes in diabetic patients with heart anomalies encompass ACE inhibitors, digoxin, diuretics,

β -blockers, Ca^{2+} antagonists, and spironolactone (7). Nonetheless, the mortality rate for diabetic cardiomyopathy still remains high, warranting novel and effective therapeutic strategies.

In this issue, Pan et al (8) extended their earlier work revealing that C66 [(2E,6E)-2,6-bis(2-(trifluoromethyl)benzylidene) cyclohexanone], a curcumin analog, reduces streptozotocin-induced hypertriglyceridemia in both serum and hearts, in addition to the significantly reduced plasma and cardiac triglyceride levels. The authors further reported that reduction in hyperlipidemia in response to C66 therapy is accompanied by improved cardiac function, inhibition of Jun NH₂-terminal kinase (JNK) signaling and cardiac inflammation in the type 1 experimental diabetic model. Pretreatment with C66 inhibited a high glucose-induced rise in proinflammatory cytokines via inactivation of nuclear factor κ B (NF- κ B). Furthermore, they showed that inhibition of JNK phosphorylation may be responsible for the beneficial effect of C66 against inflammation and apoptosis. In diabetic mice, administration of C66 or another JNK inhibitor SP600125 effectively lowered plasma and cardiac levels of tumor necrosis factor (TNF)- α , accompanied by lessened apoptosis and improved histology, including interstitial fibrosis and functional anomalies in the heart. These findings support the notion that the curcumin analog protects against diabetes and hyperglycemia-induced cardiac damage via inhibition of inflammation. More important, these findings depict a critical role for JNK activation in diabetic cardiac injury and suggest that JNK may be considered as a therapeutic target for diabetic cardiomyopathy. A recent report from the same group indicated that C66 is capable of preventing JNK activation in diabetes, leading to protection against diabetes-induced cardiac fibrosis, oxidative stress, endoplasmic reticulum (ER)

¹Center for Cardiovascular Research and Alternative Medicine, University of Wyoming College of Health Sciences, Laramie, WY

²Diabetes and Cardiovascular Center, University of Missouri School of Medicine, and VA Medical Center, Columbia, MO

Corresponding author: Jun Ren, jren@uwyo.edu, or James R. Sowers, sowersj@health.missouri.edu.

© 2014 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered.

See accompanying article, p. 3497.

stress, and cell death, along with a preserved level of the heavy metal scavenger metallothionein in the heart (9). The findings from Pan et al. (8) took it to the next level, suggesting a role of antagonism against inflammation and cytokine behind C66-offered benefits in diabetes. Inflammation with increased cytokine levels has been known to contribute to the development of diabetic cardiomyopathy (10). Both diabetes and hyperglycemia are accompanied by elevated cytokine levels, such as TNF- α and interleukin (IL)-1 β , in both cardiomyocytes and/or immunocompetent cells recruited to the heart, leading to cardiomyocyte apoptosis and ultimately cardiac dysfunction (11,12). Nonetheless, further scrutiny is needed to determine if the observed suppression of inflammation and cytokine release by C66 serve as a permissive step in the combat against diabetic cardiomyopathy (Fig. 1).

Curcumin, a natural bioactive compound derived from the root of *Curcuma longa*, has been used to treat inflammatory conditions and chronic diseases (13). In addition, curcumin is also commonly used as a coloring and flavoring additive in foods. Recent studies indicate that dietary administration of curcumin may be effective in treating conditions such as cancer, Alzheimer disease, and cystic fibrosis (14,15). With regards to the mode of action, curcumin exerts a rather diverse array of metabolic, cellular, and molecular activities (15). However, certain pharmacokinetic defects, such as low bioavailability and fast metabolism, and poor chemical stability significantly limit the clinical application of curcumin. In the past decade, this group of investigators have designed and synthesized a series of monocarbonyl analogs of curcumin including C66,

which exhibit overtly enhanced chemical stability and improved pharmacokinetic profiles in vivo (16). Their earlier work has depicted benefits of C66 in diabetic nephropathy via a mitogen-activated protein kinase (MAPK)-dependent anti-inflammatory or anti-ACE mechanism (17–19). Pan et al. (8) further demonstrated that C66, at a relatively low dosage of 5.0 mg/kg/day, protects against diabetic cardiac injury via JNK inhibition. Thus, discovery and identification of novel derivatives from natural active products may be an effective strategy for drug development in the management of diabetes complications. Although this study has revealed the therapeutic potential of C66 in the management of diabetic cardiomyopathy, more questions remain to be answered. First, can C66 be developed as a potential lipid-lowering agent, and if so, through which mechanism(s) in diabetic hyperlipidemia? Second, although JNK activation is well documented in diabetes and diabetes complications, the precise role for JNK in diabetes-induced hyperlipidemia remains unclear. The lipid uptake or transport proteins involved in JNK activation-induced change in lipid levels in the circulation and heart remain elusive. Last, but not least, diabetic cardiomyopathy was first described in conjunction with diabetic glomerulosclerosis by Rubler et al. in 1972 (20), and it was later recognized as a unique nosologic entity by the World Health Organization in 1995. Despite the large body of literature available on the pathogenesis, diagnostic, and therapeutic approaches for diabetic cardiomyopathy, its existence and etiology as a unique clinical entity continue to be debatable (21). Thus, whether C66




Figure 1—Schematic diagram depicting possible mechanism(s) involved in the beneficial effects offered by the curcumin analog C66 in diabetes/hyperglycemia-induced abnormalities in cardiac function, morphology, and cell survival. C66 inhibits JNK activation to interrupt inflammation and apoptosis during early and late phases of diabetes/hyperglycemia insult, respectively. In addition, C66 preserves levels of heavy metal scavenger metallothionein through JNK inhibition. IκB α , a regulatory protein that inhibits NF- κ B by complexing with and trapping it in the cytoplasm; P, phosphorylation; TGF- β , transforming growth factor- β .

offered a beneficial affect specific to diabetic cardiomyopathy remains to be elucidated.

Acknowledgments. The authors would like to thank Brenda Hunter from the University of Missouri School of Medicine (Columbia, MO) for her editorial assistance.

Funding. This research was supported in part by the American Diabetes Association (7-13-BS-142-BR) to J.R., National Institutes of Health grants HL-73101 and HL-107910 to J.R.S., and the Veterans Affairs Merit System 0018 (J.R.S.).

Duality of Interest. J.R.S. is a consultant for Merck Pharmaceuticals and Boehringer Ingelheim. No other potential conflicts of interest relevant to this article were reported.

References

1. Bugger H, Abel ED. Molecular mechanisms of diabetic cardiomyopathy. *Diabetologia* 2014;57:660–671
2. Boudina S, Abel ED. Diabetic cardiomyopathy revisited. *Circulation* 2007;115:3213–3223
3. Huynh K, Bernardo BC, McMullen JR, Ritchie RH. Diabetic cardiomyopathy: mechanisms and new treatment strategies targeting antioxidant signaling pathways. *Pharmacol Ther* 2014;142:375–415
4. Poirier P, Bogaty P, Garneau C, Marois L, Dumesnil JG. Diastolic dysfunction in normotensive men with well-controlled type 2 diabetes: importance of maneuvers in echocardiographic screening for preclinical diabetic cardiomyopathy. *Diabetes Care* 2001;24:5–10
5. Wold LE, Ceylan-Isik AF, Ren J. Oxidative stress and stress signaling: menace of diabetic cardiomyopathy. *Acta Pharmacol Sin* 2005;26:908–917
6. Ouyang C, You J, Xie Z. The interplay between autophagy and apoptosis in the diabetic heart. *J Mol Cell Cardiol* 2014;71:71–80
7. Pappachan JM, Varughese GI, Sriraman R, Arunagirinathan G. Diabetic cardiomyopathy: Pathophysiology, diagnostic evaluation and management. *World J Diabetes* 2013;4:177–189
8. Pan Y, Wang Y, Zhao Y, et al. Inhibition of JNK phosphorylation by a novel curcumin analog prevents high glucose–induced inflammation and apoptosis in cardiomyocytes and the development of diabetic cardiomyopathy. *Diabetes* 2014;63:3497–3511
9. Wang Y, Zhou S, Sun W, et al. Inhibition of JNK by novel curcumin analog C66 prevents diabetic cardiomyopathy with a preservation of cardiac metallothionein expression. *Am J Physiol Endocrinol Metab* 2014;306:E1239–E1247
10. Palomer X, Salvadó L, Barroso E, Vázquez-Carrera M. An overview of the crosstalk between inflammatory processes and metabolic dysregulation during diabetic cardiomyopathy. *Int J Cardiol* 2013;168:3160–3172
11. Rajesh M, Bátka S, Kechrid M, et al. Cannabinoid 1 receptor promotes cardiac dysfunction, oxidative stress, inflammation, and fibrosis in diabetic cardiomyopathy. *Diabetes* 2012;61:716–727
12. Westermann D, Van Linthout S, Dhayat S, et al. Cardioprotective and anti-inflammatory effects of interleukin converting enzyme inhibition in experimental diabetic cardiomyopathy. *Diabetes* 2007;56:1834–1841
13. Gupta SC, Patchva S, Aggarwal BB. Therapeutic roles of curcumin: lessons learned from clinical trials. *AAPS J* 2013;15:195–218
14. Lestari ML, Indrayanto G. Curcumin. In *Profiles of Drug Substances, Excipients, and Related Methodology*, Vol. 39. Brittain H, Ed. Waltham, MA, Elsevier Inc., 2014; p. 113–204
15. Zingg JM, Hasan ST, Meydani M. Molecular mechanisms of hypolipidemic effects of curcumin. *Biofactors* 2013;39:101–121
16. Liang G, Shao L, Wang Y, et al. Exploration and synthesis of curcumin analogues with improved structural stability both in vitro and in vivo as cytotoxic agents. *Bioorg Med Chem* 2009;17:2623–2631
17. Pan Y, Huang Y, Wang Z, et al. Inhibition of MAPK-mediated ACE expression by compound C66 prevents STZ-induced diabetic nephropathy. *J Cell Mol Med* 2014;18:231–241
18. Pan Y, Zhang X, Wang Y, et al. Targeting JNK by a new curcumin analog to inhibit NF- κ B-mediated expression of cell adhesion molecules attenuates renal macrophage infiltration and injury in diabetic mice. *PLoS One* 2013;8:e79084
19. Pan Y, Wang Y, Cai L, et al. Inhibition of high glucose-induced inflammatory response and macrophage infiltration by a novel curcumin derivative prevents renal injury in diabetic rats. *Br J Pharmacol* 2012;166:1169–1182
20. Rubler S, Dlugash J, Yuceoglu YZ, Kumral T, Branwood AW, Grishman A. New type of cardiomyopathy associated with diabetic glomerulosclerosis. *Am J Cardiol* 1972;30:595–602
21. Mandavia CH, Aroor AR, Demarco VG, Sowers JR. Molecular and metabolic mechanisms of cardiac dysfunction in diabetes. *Life Sci* 2013;92:601–608