

Effects of Dietary Curcumin on Glutathione S-Transferase and Malondialdehyde-DNA Adducts in Rat Liver and Colon Mucosa: Relationship with Drug Levels¹

Ricky A. Sharma,² Christopher R. Ireson,
Richard D. Verschoyle, Kirsti A. Hill,
Marion L. Williams, Chiara Leuratti,
Margaret M. Manson, Lawrence J. Marnett,
William P. Steward, and Andreas Gescher

Department of Oncology [R. A. S., K. A. H., M. L. W., W. P. S.] and MRC Centre for Mechanisms of Human Toxicity [C. R. I., R. D. V., C. L., M. M. M., A. G.], University of Leicester, Leicester LE1 9HN, United Kingdom, and Vanderbilt Cancer Center, Nashville, Tennessee 37232 [L. J. M.]

ABSTRACT

Curcumin prevents colon cancer in rodent models. It inhibits lipid peroxidation and cyclooxygenase-2 (COX-2) expression and induces glutathione S-transferase (GST) enzymes. We tested the hypothesis that 14 days of dietary curcumin (2%) affects biomarkers relevant to cancer chemoprevention in the rat. Levels of inducible COX-2, as reflected by prostaglandin E₂ production by blood leukocytes, were measured *ex vivo*. Total GST activity and adducts of malondialdehyde with DNA (M₁G), which reflect endogenous lipid peroxidation, were measured in colon mucosa, liver, and blood leukocytes. Curcumin and its metabolites were analyzed by high-performance liquid chromatography in plasma, and its pharmacokinetics were compared following a diet containing 2% curcumin *versus* intragastric (i.g.) administration of curcumin suspended in an amphiphilic solvent. The curcumin diet did not alter any of the markers in the blood but increased hepatic GST by 16% and decreased colon M₁G levels by 36% when compared with controls. Administration of carbon tetrachloride during the treatment period increased colon M₁G levels, and this increase was prevented by dietary curcumin. Dietary curcumin yielded low drug levels in the plasma, between 0 and 12 nM, whereas tissue concentrations of curcumin in liver and colon mucosa were 0.1–0.9 nmol/g and 0.2–1.8 μmol/g, re-

spectively. In comparison with dietary administration, suspended curcumin given i.g. resulted in more curcumin in the plasma but much less in the colon mucosa. The results show that curcumin mixed with the diet achieves drug levels in the colon and liver sufficient to explain the pharmacological activities observed and suggest that this mode of administration may be preferable for the chemoprevention of colon cancer.

INTRODUCTION

Cancer mortality rates in the developed world have risen throughout most of this century, and it is already the leading cause of death in some Western countries (1, 2). This observation has engendered much research activity aimed at the identification of cancer chemopreventive agents, especially substances derived from the diet (3). Turmeric, the dried ground rhizome of the perennial herb *Curcuma longa*, is an example of one such agent. This spice is consumed in the diet in quantities up to 4 g adult/day in some countries (4), which also appear to have low incidence rates of colorectal cancer (1). Turmeric contains curcuminoids and essential oils. Curcumin (diferuloylmethane), a low molecular weight polyphenol and the major curcuminoid in the plant, is regarded as the constituent with the highest biological activity. Curcumin has been shown to inhibit tumor formation in the skin, forestomach, duodenum, and colon of mice and in the tongue, colon, mammary glands, and sebaceous glands of rats (5). Mechanisms by which curcumin causes cancer chemoprevention are thought to involve antioxidation (6), inhibition of kinases (7), interference with the activity of transcription factors such as nuclear factor-κB and activator protein-1 (8), and suppression of expression of the enzyme COX-2³ (9).

In a program of preclinical work designed to help optimize the clinical evaluation of curcumin as a colon cancer chemopreventive agent, we tested the hypothesis that curcumin in the diet alters biomarkers of its chemopreventive efficacy. Three biomarkers were selected to represent possible mechanisms of the chemopreventive activity of curcumin *in vivo*. These were total GST activity, DNA adducts formed by MDA, and inducible COX-2 expression as reflected by PGE₂ production. Induction of GSTs, phase II enzymes that detoxify certain carcinogens, is regarded as a potential mechanism of blockade of the early stages of carcinogenesis (3). Such induction is a property

Received 12/8/00; revised 1/31/01; accepted 2/5/01.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked advertisement in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

¹ Supported in part by a Leicester Royal Infirmary Research Fellowship (to R. A. S.), United Kingdom Ministry for Agriculture, Fisheries, and Food Contract FS1735 (to C. L.), and NIH Grant CA-77839 (to L. J. M.).

² To whom requests for reprints should be addressed, at Oncology Department, Hodgkin Building, University of Leicester, Leicester LE1 9HN, United Kingdom. Phone: 44-116-252-5541; Fax: 44-116-252-5616; E-mail: ras20@le.ac.uk.

³ The abbreviations used are: COX, cyclooxygenase; GST, glutathione S-transferase; MDA, malondialdehyde; PG, prostaglandin; M₁G, malondialdehyde-deoxyguanosine adduct; HPLC, high-performance liquid chromatography; LPS, lipopolysaccharide; CDNB, 1-chloro-2,4-dinitrobenzene as substrate; i.g., intragastric.

of several agents with chemopreventive activity in preclinical models, including curcumin (10) and the broccoli constituent sulforaphane (11). MDA is a naturally occurring product of lipid peroxidation (12), which is also formed during prostaglandin biosynthesis via COX (13), two enzymatic processes that have been implicated in the pathogenesis of a number of cancers, especially colon cancer (14–16). PGE₂ is a product of COX-2, the isoenzyme of COX induced during infection, inflammation, and malignant transformation. Inhibition of COX-2 is thought to be an important mechanism of chemoprevention, exhibited by nonsteroidal anti-inflammatory drugs (14) and dietary agents such as curcumin (9). MDA reacts with DNA under physiological conditions to form adducts, predominantly with deoxyguanosine (M₁G). Accumulated DNA damage may be important in the etiology of many cancers, and such damage may be reflected by exocyclic DNA adducts such as M₁G (17). M₁G has been implicated in the induction of G→T transversions by MDA and is considered a potentially useful “dosimeter” of MDA-induced DNA modification (18). M₁G levels have been described in malignant and nonmalignant human colon cells *in vitro*⁴ and in leukocytes, pancreas, breast, and liver in healthy volunteers (12); indeed they are comparable with levels of DNA adducts formed by exogenous carcinogens such as polycyclic hydrocarbons.

The ability of curcumin to prevent malignancies in the colorectal tract in rodents has been particularly well documented (7, 19, 20). It has also been shown to inhibit lipid peroxidation (21) and carbon tetrachloride-mediated hepatotoxicity (22, 23). We therefore aimed to test the hypothesis that dietary curcumin affects GST and M₁G levels in the intestinal mucosa and prevents increases in M₁G levels when lipid peroxidation is induced in this tissue by CCl₄. For comparison, levels of GST and M₁G were also investigated in liver and blood leukocytes. Because COX-2 is not expressed significantly in normal tissues without provocation, an *ex vivo* model was used to study its inducibility in peripheral blood indirectly. To be able to rationalize potential pharmacodynamic changes and interpret them in terms of efficacious drug levels, curcumin and its glucuronide or sulfate conjugation products were determined in intestinal mucosa, plasma, and liver. Studies in rodents suggest that curcumin has poor systemic bioavailability when given *p.o.* and is cleared rapidly from the plasma when administered *i.v.* (24), but overall the information on its pharmacokinetic behavior is scarce. Whereas in cancer intervention studies using curcumin it has generally been administered admixed in the diet, studies of its pharmacokinetics and metabolism have used *i.g.* gavage or non-oral routes of administration. In the light of the difficult pharmaceutical properties of curcumin, *i.e.*, its lipophilicity and susceptibility to rapid metabolism, we also tested the hypothesis that drug and metabolite levels after *i.g.* gavage are representative of those following curcumin ingestion in the diet.

⁴R. A. Sharma, A. Gescher, J. P. Plataras, C. Leuratti, R. Singh, L. J. Marnett, W. P. Steward, and S. M. Plummer. Role of cyclooxygenase-2 in modulation of malondialdehyde levels and its deoxyguanosine adduct in human colon cells. *Carcinogenesis* (Lond.), submitted for publication.

MATERIALS AND METHODS

Reagents. Curcumin was purchased from Apin Chemicals Ltd. (Abingdon, United Kingdom). Analysis by HPLC/mass spectrometry established that the material contained 91% curcumin and 9% desmethoxycurcumin. CCl₄, corn oil, glutathione, CDNB, glycerol formal (consisting of 60% 5-hydroxy-1,3-dioxane and 40% 4-hydroxymethyl-1,3-dioxalane) and Cremophore were purchased from Sigma Chemical Co. (Poole, United Kingdom). Murine M₁G monoclonal antibody D10A1 was prepared as described previously (25). Antimurine horseradish peroxidase antibody was purchased from Dako (Ely, United Kingdom). M₁G standards were synthesized and characterized as described previously (26).

Treatments. Female F344 rats (6-weeks of age; 160–180 g), obtained from Charles River UK Ltd. (Margate, United Kingdom) had access to water and a standard RMI diet (Special Diet Services) *ad libitum*. Rats were kept in groups of four per cage at 20°C on a 12-h light/dark cycle. For dietary administration, curcumin and corn oil were mixed with the diet to furnish 2% for each. This level of curcumin in the diet, which rats received for 14 days, has been shown previously to prevent gastrointestinal tract tumors in several rodent models of chemically induced carcinogenesis (5, 10). Control animals received a diet containing corn oil. CCl₄ was dissolved in corn oil (2 ml/kg) and administered by *i.g.* gavage at 0.5 ml/kg on day 10 after commencement of dietary curcumin intake. This dose of CCl₄ has been shown previously to raise liver M₁G levels (18). Control animals received the equivalent dose of corn oil.

In the pharmacokinetic study, administration of dietary curcumin was compared with that of an *i.g.* bolus. For administration by gavage, curcumin was suspended and partially dissolved in a mixture of glycerol formal:cremophore:water (5:2:2), which in preliminary experiments was found to yield a suitable formulation in terms of acceptable viscosity and stability of suspension. This formulation was administered at 500 mg curcumin/kg by *i.g.* gavage once only or daily for 7 consecutive days. Control animals received the excipient mixture only. Tissue samples were collected 30 min postdose, a time point at which in preliminary experiments curcumin levels had been found to be maximal. For an optimal comparison of this administration mode with dietary curcumin, we exploited the fact that feeding habits of rats are subject to a diurnal cycle with two eating peaks, one of which occurs at around 8:00 p.m. (27). Rats were deprived of food for 6 h, commencing at 1:00 p.m. They then received the curcumin-containing diet for 3 h, coinciding with the eating peak. Subsequently, food was withdrawn for ~30 min, after which blood and tissue samples were collected. This “starvation-refeeding” protocol was performed either in unpretreated animals or in rats that had received the curcumin diet continually for 7 days.

At the end of the feeding period, or subsequent to *i.g.* administration, rats were subjected to terminal anesthesia (halothane/nitrous oxide), and blood was removed by cardiac puncture. Blood was placed in heparinized tubes and plasma obtained by centrifugation (1100 × *g* at 4°C for 25 min). The large intestine was flushed with PBS and dissected out. Cytosol was prepared by standard procedures (28), and blood leukocytes were isolated using Ficoll-Paque Plus (Amersham Pharmacia

Biotech, Buckinghamshire, United Kingdom). Histological examination ensured accurate scraping of colonic mucosa only. Blood and tissue samples were protected from light, frozen in liquid nitrogen, and stored at -80°C until pharmacodynamic and pharmacokinetic analyses were performed.

Pharmacodynamic Analyses. Total GST activity of cytosol samples was measured spectrophotometrically using glutathione and CDNB (29). Results were corrected for protein levels using the Bio-Rad protein assay (Bio-Rad, Hemel Hempstead, United Kingdom). Extraction of genomic DNA and analysis of M₁G adduct levels by immunoslot blot was performed as described previously (30). Discrepancies in the amount of DNA in each slot were corrected by staining the nitrocellulose filter with propidium iodide and performing UV light densitometry. The detection limit for M₁G was five adducts per 10^8 nucleotides. The model for assessing inducibility of COX-2 *ex vivo* was based on published methods and has been well established in clinical studies as well as preclinical models (31, 32). Aliquots (0.5 ml) of fresh blood were incubated with acetylsalicylic acid (200 μM) for 30 min at 37°C to inactivate platelet COX-1 irreversibly. LPS (10 $\mu\text{g}/\text{ml}$) was added to half the samples and mixed well, and samples were reincubated for 24 h. Plasma was separated by centrifugation and stored at -80°C . PGE₂ in these plasma samples was measured by competitive enzyme immunoassay (Cayman Chemicals, Ann Arbor, MI), with a detection limit of ~ 30 pg/ml plasma.

Analysis of Curcumin and Its Conjugates. Curcumin and its metabolites were measured as described before (24). Plasma and tissues were extracted with twice the volume of ethyl acetate, and the organic layer was evaporated under nitrogen. Extraction efficiencies from plasma for curcumin, curcumin glucuronide, and curcumin sulfate determined by HPLC (see below) at 0.1 $\mu\text{g}/\text{ml}$ were 92 ± 7 , 45 ± 10 , and $49 \pm 9\%$ (mean \pm SD, $n = 6$), respectively; the extraction efficiencies at 40 $\mu\text{g}/\text{ml}$ were very similar to these values. Extraction efficiencies from liver and colon mucosal scrapings were not significantly different from those obtained in plasma. The reversed-phase HPLC method for detection and quantitation of curcumin and its conjugates used a Symmetry Shield RP 18 column (150 \times 3.9 mm; particle size, 5 μm ; Waters) and a Varian Prostar (230 model) solvent delivery system coupled to a UV-visible detector (310 model) and autosampler (model 410). Detection of curcumin, curcumin sulfate, and curcumin glucuronide was achieved at 420 nm. For the detection of products of curcumin reduction, the detector was switched to 280 nm. 5,10,15,20-Tetra-(*m*-hydroxyphenyl)-chlorin was used as an internal standard. Samples were reconstituted in acetonitrile:water (1:1), and the injection volume was 50 μl . A linear gradient of 5–45% acetonitrile in 0.01% ammonium acetate (pH 4.5) was used for 30 min, followed by an increase over 20 min to 95% acetonitrile (flow rate, 1 ml/min). The limits of detection for curcumin, curcumin glucuronide, and curcumin sulfate were 5 pmol/ml plasma or 25 pmol/g tissue, and the limits of quantitation were near 10 pmol/ml plasma and 50 pmol/g tissue. Quantitation of the conjugates was based on calibration curves obtained for curcumin, because we had established previously that the specific absorption at 420 nm was almost identical for the three compounds (24). The quantitative method for curcumin was validated using a 0.1 and 40 $\mu\text{g}/\text{ml}$ solution yielding

Table 1 Effect of dietary curcumin^a on GST levels in rat liver, colon mucosa, and blood lymphocytes

Tissue	Control	Curcumin-fed
Liver	634 \pm 66 ^b	860 \pm 73 ^c
Colon mucosa	84 \pm 13	69 \pm 16
Blood lymphocytes	20 \pm 4	16 \pm 4

^a Animals were exposed to curcumin (2%) in the diet for 14 days.

^b Values, expressed as nmol/min/mg protein using CDNB as substrate, are the mean \pm SD of eight separate animals. For experimental details see "Materials and Methods."

^c The difference between control and treated animals is significant ($P < 0.01$, by ANOVA).

intra-day and inter-day coefficients of variation of 17 and 13%, and 16 and 19%, respectively ($n = 4$).

Statistical Evaluation. Results were subjected to ANOVA and Spearman's test of rank correlation, using Excel and Minitab (both Windows 1997) software packages. *Post hoc* Fisher's Least Significant Difference Test was performed.

RESULTS

Pharmacodynamic Effects of Dietary Curcumin. Rats were fed a diet supplemented with 2% curcumin for 2 weeks. On the basis of food consumption, the approximate daily dose ingested was 1.2 g curcumin/kg. Dietary curcumin at this dose did not affect the animals' body weight gain. Levels of GST activity in the liver of curcumin-fed rats were elevated by 36% over those in control animals (Table 1). GST levels in colon mucosa and lymphocytes were marginally decreased by curcumin; however, this difference was not significant. Levels of M₁G adducts in colon mucosa of rats which received curcumin were moderately, but significantly, lower than those in control rats (Fig. 1). M₁G adduct levels in leukocytes and liver of curcumin-fed rats were also slightly decreased as compared with controls, but these differences were not significant. Treatment of rats with CCl₄ increased levels of M₁G adducts in liver and colon mucosa by 49 and 25%, respectively, over control values (Fig. 1). Dietary curcumin attenuated this increase in the liver and completely prevented it in the colon.

To study whether dietary curcumin led to altered COX-2 expression in blood cells, the concentration of PGE₂ in plasma was determined after induction with LPS added *in vitro*. LPS increased blood PGE₂ levels ~ 3 -fold, but curcumin feeding failed to interfere with this increase (result not shown).

Levels of Curcumin after Dietary Administration. Plasma, colon mucosa, and liver from rats that had received curcumin in their diet as described above for 2 weeks were analyzed for the presence of curcuminoids. Curcumin and its metabolites, curcumin glucuronide and curcumin sulfate, could not be detected in the plasma obtained by cardiac puncture or from the hepatic portal vein. Levels of curcumin were 1.8 ± 0.8 $\mu\text{mol}/\text{g}$ tissue in the colon mucosa and 0.8 ± 0.3 nmol/g in the liver. Curcumin was also present in the feces (8.6 ± 0.6 $\mu\text{mol}/\text{g}$ dried feces). Curcumin glucuronide or curcumin sulfate was not detected in either tissues or feces.

Pharmacokinetic Comparison of Modes of Administration. To compare the availability of curcumin admixed with the diet or suspended in an amphiphilic solvent and adminis-




Fig. 1 Effect of dietary curcumin on M₁G adduct levels in liver, colon mucosa, and blood leukocytes. □, control animals; ■ and ▨, animals that received dietary curcumin (2%) for 14 days; ▩ and ▪, animals that received carbon tetrachloride (0.5 ml/kg) via the i.g. route on the 10th day of the study diet. Values are means of eight animals; bars, SD. * and **, the difference between control and curcumin-treated animals (*, $P < 0.01$, by ANOVA) or between animals that received carbon tetrachloride either with or without curcumin (**, $P < 0.005$, by ANOVA) is significant. For experimental details, see "Materials and Methods."

Table 2 Curcumin and curcumin conjugates in the plasma of rats that received curcumin in the diet or by the i.g. route

Dosing schedule ^a	Plasma levels (pmol/ml)		
	Curcumin	Curcumin glucuronide	Curcumin sulfate
Short feeding	<10 ^b	ND ^c	ND
Single i.g.	30 ± 9	56 ± 36	134 ± 100
7-day feeding	12 ± 5	ND	ND
Repeated i.g.	65 ± 28	638 ± 146	360 ± 160

^a Rats received curcumin as an i.g. bolus (500 mg/kg) once ("Single i.g.") or daily on 7 consecutive days ("Repeated i.g."), or with the diet (2%) for 3 h after a period of starvation (6 h) once ("Short feeding") or after unlimited access to curcumin containing diet for 7 days ("7-day feeding"). The dose, as calculated by weighing food removal, was between 200 and 400 mg/kg. For details of curcumin administration and measurement of curcumin and metabolites, see "Materials and Methods."

^b Values are mean ± SD from three to four animals.

^c ND, not detectable.

tered by i.g. bolus, we studied drug levels after short-term feeding (2% in the diet for 3 h) or single i.g. bolus (500 mg/kg) and after continual feeding or repeated bolus administration for 7 days. To render the dietary dose ingested and the sample timing comparable with those associated with the i.g. bolus, we exploited the "starvation-refeeding" protocol described in "Materials and Methods." The dose ingested with the diet during the refeeding period, as adjudged by the amount of food removed from the hoppers, was between 200 and 400 mg/kg, *i.e.*, between 40 and 80% of, and thus comparable with, the i.g. dose. Tables 2 and 3 show that the levels of curcumin and its conjugates differed substantially, depending on the mode of administration. Plasma levels of curcumin were approximately three to five times higher after the i.g. bolus than after dietary ingestion, and curcumin conjugates were observed in the plasma only when the agent was given via the i.g. route (Table 2). Repeated

Table 3 Curcumin levels in colon mucosa of rats that received curcumin in the diet or by the i.g. route

Dosing schedule ^a	Curcumin levels (nmol/g tissue)
Short feeding	279 ± 295
Single i.g.	1.7 ± 0.9 ^b
7-day feeding	482 ± 412
Repeated i.g.	18 ± 24

^a Rats received curcumin as an i.g. bolus (500 mg/kg) once ("Single i.g.") or daily for 7 consecutive days ("Repeated i.g."), or with the diet (2%) for 3 h after a period (6 h) of starvation ("Short feeding"), or after a starvation-refeeding regimen after unlimited access to a curcumin-containing diet for 7 days ("7-day feeding"). The dose of curcumin when given with the diet, as calculated by weighing food removal, was between 200 and 400 mg/kg. For details of curcumin administration and its measurement, see "Materials and Methods."

^b Values are mean ± SD from three to four animals.

i.g. administration for 7 days led to an increase in plasma levels of curcumin and its conjugates as compared with single i.g. administration. Curcumin levels were doubled, and levels of curcumin sulfate and curcumin glucuronide were 2.7 and 11 times higher than those seen after single i.g. administration. In contrast, plasma levels of curcumin after its consumption in the diet for a week, at the end of which rats were subjected to the starvation-refeeding regime, did not differ substantially from those measured after short-term dietary consumption. The variation between animals in colon mucosal curcumin levels (Table 3) after dietary consumption was very high, probably reflecting considerable differences in food intake between individual animals during the 3-h period of refeeding. Curcumin levels in the colon mucosa after short-term dietary consumption were 164 times higher than those seen after single i.g. bolus; after dietary consumption for 1 week, they were 27 times higher than after the last of seven daily i.g. bolus doses (Table 3). Liver levels of curcumin were ~0.1 nmol/g, irrespective of route of administration, and there was little difference in liver levels between single or repeated administration via the diet or gavage.

DISCUSSION

The results of this study allow two conclusions that may help to optimize clinical trials of curcumin as a cancer chemopreventive agent: (a) the bioavailability of curcumin in blood and tissues is dramatically affected by the way in which it is p.o. administered; and (b) colon mucosa and liver are pharmacological targets of dietary curcumin.

The first conclusion is based on the observed differences in levels of curcumin and its conjugates in plasma and tissues after the two administration modes. Dietary curcumin elicited concentrations of the drug in the colon mucosa of between 0.3 and 1.8 μmol/g, whereas plasma levels were around the limit of detection. Curcumin suspended in a solvent mixture and given by i.g. bolus furnished levels of drug in the plasma that were 3–6-fold higher than those seen after dietary administration. Curcumin metabolites were detectable in the plasma only after i.g. administration. Conversely, colon mucosal levels of curcumin after i.g. bolus were only a fraction of those observed after dietary administration.

Curcumin exerts its effects on growth, COX-2 expression,

and transcription factor activity in cells *in vitro* at concentrations of 5 μM or above (9). The colon mucosal level of curcumin after 14 days feeding observed in this study, 1.8 $\mu\text{mol/g}$, is more than 300-fold higher than the minimal concentration shown to be active *in vitro*. This result demonstrates that dietary administration of curcumin can produce pharmacologically relevant drug concentrations in colon mucosa. Although the dose used in this study, 2% in the diet, has been used frequently in rodent intervention studies (5, 10, 33), it is at least 10 times higher than the highest estimated daily human intake of curcumin as a dietary constituent (4) and does not reflect normal dietary use. In recent intervention studies, dietary levels of 0.2 and 0.1% curcumin have been demonstrated to protect rodents from colon cancer induced by azoxymethane (7, 19, 20) and to reduce polyp numbers in the Min mouse (34), a model of the genetic defect found in human familial adenomatous polyposis. Our results therefore intimate that dietary curcumin at doses considerably lower than those used here might yield pharmacologically efficacious levels in the colon mucosa and perhaps also in the liver.

Intriguingly, curcumin glucuronide and curcumin sulfate were not detected in plasma or tissues after its administration in the diet, whereas they were the major drug-derived species present in the plasma after i.g. administration. In a recent report, a small amount of a curcumin reduction product, tetrahydrocurcumin, was found in plasma samples from rats fed a curcumin-containing diet (1%) subsequent to treatment of plasma with enzymes that hydrolyze xenobiotic glucuronides and sulfates (35). In contrast, products of curcumin reduction were not unequivocally detected in plasma after dietary curcumin given for 14 days in the experiments described here, although hexahydrocurcumin and hexahydrocurcuminol were identified by mass spectrometry in the bile.⁵ Overall, this part of the study suggests that the pharmacokinetic behavior of curcumin after administration of an i.g. bolus of curcumin in suspension is clearly unrepresentative of that of curcumin mixed into the diet. This finding is important because, although in intervention studies curcumin is generally given as a constituent of the diet, studies of its pharmacokinetics and metabolism have been performed mostly with the drug formulated in suspension, using DMSO, aqueous carboxymethyl cellulose solution, or arachis oil as solvents, and administered as an i.g. bolus or via the i.p. route (36–39). The results presented here suggest that should oral curcumin be advocated in the chemoprevention of malignancies remote from the liver or gastrointestinal tract, improvement of its oral bioavailability might be necessary, perhaps by formulating it as a solution.

Our second conclusion is borne out by the findings that dietary curcumin elevated hepatic GST levels, reduced colon mucosal M₁G adduct levels, and decreased the elevation of M₁G adduct levels elicited by a powerful lipid peroxidative stimulus in the liver and colon mucosa. These potentially beneficial effects correlated with curcumin levels of 1.8 $\mu\text{mol/g}$ in the colon mucosa and 0.8 nmol/g in liver. The high concentrations of curcumin in the colon mucosa described here are consistent with the outcome of intervention studies in preclinical rodent

models of colon cancer (7, 19, 20, 34). Although the decrease in M₁G levels in colon mucosa was modest, it underlines in principle that curcumin supplementation of the diet can achieve drug concentrations sufficient to decrease levels of DNA adducts formed as a corollary of lipid peroxidation. Curcumin glucuronide and curcumin sulfate were not found in liver or colon mucosa after dietary administration of curcumin, strongly suggesting that parent curcumin rather than either conjugate affects GST activities and M₁G adduct levels *in vivo*.

The levels of M₁G adducts in rat liver measured here were approximately three times higher than those described previously in Sprague Dawley rats and are more analogous to background levels found in normal human liver (18). Differences may be related to the age, sex, and strain of the animals used, or fat composition of the diets used. M₁G adduct levels in rat leukocytes and colon mucosa have not been documented previously. CCl₄ has been shown to increase MDA and M₁G adduct levels in rat liver (18), and increases in hepatic MDA levels have been attenuated by feeding rats 100 mg/kg curcumin for 4 weeks (23). The results described here extend these findings, because they show that CCl₄ also engenders M₁G adducts in the colon mucosa, and that this increase can be prevented by dietary curcumin. These findings are pertinent in the light of a recent study in which oxidative DNA damage linked to lipid peroxidation was detected in biopsies from normal human colon (40). The M₁G-lowering effect of curcumin provides a tentative rationale for the regular use of dietary curcumin in the protection of the colon mucosa against oxidative damage, perhaps in pre-malignant conditions such as ulcerative colitis.

Liver GST activity was raised over controls after curcumin ingestion in the diet. The effects of curcumin on GST and its expression are complex and may involve competitive enzyme inhibition (41) as well as indirect enzyme induction (42). Our finding that 14 days of dietary curcumin induces GST activity in the liver is consistent with an earlier study of the same dose of curcumin in diet and water fed to 8-week-old mice (10). Similarly, mice that received curcumin dissolved in aqueous sodium carboxymethylcellulose via i.g. gavage for 15 days at a dose approximately one-fifth of that used here were found to have significantly higher GST levels in liver compared with controls (41). In contrast, in a more recent study of 8-week-old Sprague Dawley rats, curcumin dissolved in corn oil given daily for 14 days via the i.g. route at various doses failed to induce hepatic GST at doses >6% of the dietary dose used here (43). This discrepancy may relate to differences between the studies in rodent species and strain, age of the animals, dose of curcumin, and its route of administration. Compatible with earlier rodent studies of dietary curcumin (10, 44), we failed to detect any significant alteration in GST activity in the colonic mucosa.

In cells *in vitro*, we have shown previously that curcumin inhibits COX-2 expression by a mechanism involving interference with nuclear factor- κB activation and inhibition of the I κB kinase complex (9). In the study described here, PGE₂ production induced *ex vivo* as an indicator of leukocyte COX-2 activity was not affected by dietary curcumin. The fact that curcumin administered in the diet did not affect COX-2 inducibility, GST activity, or M₁G levels in rat blood leukocytes is consistent with the finding that this mode of administration furnished extremely low levels of parent curcumin in the plasma.

⁵ Unpublished result.

In conclusion, the results presented above show that dietary administration of curcumin to rats produces pharmacologically active levels of unmetabolized curcumin in the colon mucosa and liver, capable of decreasing M₁G levels and elevating GST activity. These effects may contribute to cancer chemoprevention. The results also suggest that dietary admixture may be the preferable mode of administration for curcumin in the chemoprevention of colon cancer. The chemopreventive efficacy of oral curcumin in the colorectum and the liver merits clinical evaluation.

ACKNOWLEDGMENTS

We thank Samuel Oustric, Simon Plummer, Raj Singh, Eliot Deag, and Don Jones for technical assistance and Clare Winfield and Sue Spriggs for helping to prepare the manuscript.

REFERENCES

- Greenlee, R. T., Murray, R., Bolden, S., and Wingo, P. A. Cancer statistics, 2000. *CA Cancer J. Clin.*, *50*: 7–33, 2000.
- Tattersall, M. H. N., and Thomas, H. Recent advances: oncology. *Br. Med. J.*, *318*: 445–448, 1999.
- Sharma, R. A., Manson, M. M., Gescher, A., and Steward, W. P. Colorectal cancer chemoprevention: biochemical targets and clinical development of promising agents. *Eur. J. Cancer*, *37*: 12–22, 2001.
- Sambaiah, K., Ratankumar, S., Kamanna, V. S., Satyanarayana, M. N., and Rao, M. V. L. Influence of turmeric and curcumin on growth, blood constituents, and serum enzymes in rats. *J. Fed. Sci. Technol. (Mysore)*, *19*: 187–190, 1982.
- Kelloff, G. J., Crowell, J. A., Hawk, E. T., Steel, V. E., Lubet, R. A., Boone, C. W., Covey, J. M., Doody, L. A., Omenn, G. S., Greenwald, P., Hong, W. K., Parkinson, D. R., Bagheri, D., Baxter, G. T., Blunden, M., Doeltz, M. K., Eisenhauer, K. M., Johnson, K., Knapp, G. G., Longfellow, D. G., Malone, W. F., Nayfield, S. G., Seifried, H. E., Swall, L. M., and Sigman, C. C. Strategy and planning for chemopreventive drug development: Clinical Development Plans II. *J. Cell. Biochem.*, *26S*: 54–71, 1996.
- Jovanovic, S. V., Steenken, S., Boone, C. W., and Simic, M. G. H-atom transfer is a preferred antioxidant mechanism of curcumin. *J. Am. Chem. Soc.*, *121*: 9677–9681, 1999.
- Rao, C. V., Rivenson, A., Simi, B., and Reddy, B. S. Chemoprevention of colon carcinogenesis by dietary curcumin, a naturally occurring plant phenolic compound. *Cancer Res.*, *55*: 259–266, 1995.
- Huang, T. S., Lee, S. C., and Lin, J. K. Suppression of c-jun/AP-1 activation by an inhibitor of tumor promotion in mouse fibroblast cells. *Proc. Natl. Acad. Sci. USA*, *88*: 5292–5296, 1991.
- Plummer, S. M., Holloway, K. A., Manson, M. M., Munks, R. J. L., Kaptein, A., Farrow, S., and Howells, L. Inhibition of cyclooxygenase 2 expression in colon cells by the chemopreventive agent curcumin involves inhibition of NF- κ B activation via the NIK/IKK signaling complex. *Oncogene*, *18*: 6013–6020, 1999.
- Singh, S. V., Hu, X., Srivastava, S. K., Singh, M., Xia, H., Orchard, J. L., and Zaren, H. A. Mechanism of inhibition of benzo[a]pyrene-induced forestomach cancer in mice by dietary curcumin. *Carcinogenesis (Lond.)*, *19*: 1357–1360, 1998.
- Zhang, Y., Kensler, T. W., Cho, C. G., Posner, G. H., and Talalay, P. Anticarcinogenic activities of sulforaphane and structurally related synthetic nortranylyl isothiocyanates. *Proc. Natl. Acad. Sci. USA*, *91*: 3147–3150, 1994.
- Marnett, L. J. Lipid peroxidation–DNA damage by malondialdehyde. *Mutat. Res.*, *424*: 83–95, 1999.
- Diczfalusy, U., Falardeau, P., and Hammarstrom, S. Conversion of prostaglandin endoperoxides to C17-hydroxyacids catalysed by human platelet thromboxane synthase. *FEBS Lett.*, *84*: 271–274, 1977.
- Taketo, M. N. Cyclooxygenase-2 inhibitors in tumorigenesis (part II). *J. Natl. Cancer Inst.*, *90*: 1609–1620, 1998.
- Hendrickse, C. W., Kelly, R. W., Radley, S., Donovan, I. A., Keighley, M. R. B., and Neoptolemos, J. P. Lipid peroxidation and prostaglandins in colorectal cancer. *Br. J. Surg.*, *81*: 1219–1223, 1994.
- Wang, M. Y., Dhingra, K., Hittelman, W. N., Liehr, J. G., de Andrade, M., and Li, D. H. Lipid peroxidation-induced putative malondialdehyde–DNA adducts in human breast tissues. *Cancer Epidemiol. Biomark. Prev.*, *5*: 705–710, 1996.
- Marnett, L. J. Oxyradicals and DNA damage. *Carcinogenesis (Lond.)*, *21*: 361–370, 2000.
- Chaudhary, A. K., Nokubo, M., Reddy, G. R., Yeola, S. N., Morrow, J. D., Blair, I. A., and Marnett, L. J. Detection of endogenous malondialdehyde–deoxyguanosine adducts in human liver. *Science (Wash. DC)*, *265*: 1580–1582, 1994.
- Samaha, H. S., Kelloff, G. J., Steele, V., Rao, C. V., and Reddy, B. S. Modulation of apoptosis by sulindac, curcumin, phenylethyl-3-methylcaffeate, and 6-phenylhexyl isothiocyanate: apoptotic index as a biomarker in colon cancer chemoprevention and promotion. *Cancer Res.*, *57*: 1301–1305, 1997.
- Kawamori, T., Lubet, R., Steele, V. E., Kelloff, G. J., Kaskey, R. B., Rao, C. V., and Reddy, B. S. Chemopreventive effect of curcumin, a naturally occurring anti-inflammatory agent, during the promotion/progression stages of colon cancer. *Cancer Res.*, *59*: 597–601, 1999.
- Ramirez Bosca, A., Soler, A., Gutierrez, M. A. C., Alvarez, J. L., and Almagro, E. Q. Antioxidant *Curcuma* extracts decrease the blood lipid levels of human subjects. *Age*, *18*: 167–169, 1995.
- Subramanian, L., and Selvam, R. Prevention of CCl₄-induced hepatotoxicity by aqueous extract of turmeric. *Nutr. Res.*, *19*: 429–441, 1999.
- Park, E. J., Jeon, C. H., Ko, G., Kim, J., and Sohn, D. H. Protective effect of curcumin in rat liver injury induced by carbon tetrachloride. *J. Pharm. Pharmacol.*, *52*: 437–440, 2000.
- Ireson, C. R., Orr, S., Jones, D. L., Verschoyle, R., Lim, C. K., Luo, J. L., Howells, L., Plummer, S. P., Jukes, R., Williams, M., Steward, W. P., and Gescher, A. Characterization of metabolites of the chemopreventive agent curcumin in humans and rat hepatocytes and in rat plasma and evaluation of their ability to inhibit cyclooxygenase-2 expression. *Cancer Res.*, *61*: 1058–1064, 2001.
- Sevilla, C. L., Mahle, N. H., Eliezer, N., Uzieblo, A., O'Hara, S. M., Nokubo, M., Miller, R., Rouzer, C. A., and Marnett, L. J. Development of monoclonal antibodies to the malondialdehyde–deoxyguanosine adduct, pyrimidopurinone. *Chem. Res. Toxicol.*, *10*: 172–180, 1997.
- Chapeau, M. C., and Marnett, L. J. Enzymatic synthesis of purine deoxynucleoside adducts. *Chem. Res. Toxicol.*, *4*: 636–638, 1991.
- Vermeulen, J. K., De Vries, A., Schlingmann, F., and Remie, R. Food deprivation: common sense or nonsense? *Anim. Technol.*, *48*: 45–54, 1997.
- Manson, M. M., Ball, H. W. L., Judah, D. J., Williamson, G., and Neal, G. E. Mechanism of action of dietary chemoprotective agents in rat liver: induction of phase I and II drug-metabolizing enzymes and aflatoxin B1 metabolism. *Carcinogenesis (Lond.)*, *18*: 1729–1738, 1997.
- Habig, W. H., Pabst, M. J., and Jakoby, W. B. Glutathione S-transferases. *J. Biol. Chem.*, *249*: 7130–7139, 1974.
- Lauranti, C., Singh, R., Lagneau, C., Farmer, P. B., Marnett, L. J., and Shuker, D. E. G. Determination of malondialdehyde-induced DNA damage in human tissues using an immunoslot blot assay. *Carcinogenesis (Lond.)*, *19*: 1919–1924, 1998.
- Patrignani, P., Panara, M. R., Greco, A., Fusco, O., Natoli, C., Iacobelli, S., Cipollone, F., Ganci, A., Crémion, C., Maclouf, J., and Patrono, C. Biochemical and pharmacological characterization of the cyclooxygenase activity of human blood prostaglandin endoperoxide synthases. *J. Pharmacol. Exp. Ther.*, *271*: 1705–1712, 1994.
- Warner, T. D., Giuliano, F., Vojnovic, I., Bukasa, A., Mitchell, J. A., and Vane, J. R. Nonsteroidal drug selectivities for cyclooxygenase-1 rather than cyclooxygenase-2 are associated with human gastrointestinal toxicity: a full *in vitro* analysis. *Proc. Natl. Acad. Sci. USA*, *96*: 7563–7568, 1999.

33. Huang, M. T., Newmark, H. L., and Frenkel, K. Inhibitory effects of curcumin on tumorigenesis in mice. *J. Cell. Biochem.*, *67*: 26–34, 1994.
34. Mahmoud, N. N., Carothers, A. M., Grunberger, D., Bilinski, R. T., Churchill, M. R., Martucci, C., Newmark, H. L., and Bertagnolli M. M. Plant phenolics decrease intestinal tumors in an animal model of familial adenomatous polyposis. *Carcinogenesis (Lond.)*, *21*: 921–927, 2000.
35. Inano, H., Onoda, M., Inafuku, N., Kubota, M., Kamada, Y., Osawa, T., Kobayashi, H., and Wakabayashi, K. Potent preventive action of curcumin on radiation-induced initiation of mammary tumorigenesis in rats. *Carcinogenesis (Lond.)*, *21*: 1835–1841, 2000.
36. Holder, G. M., Plummer, J. L., and Ryan, A. J. The metabolism and excretion of curcumin [1,7-bis-(4-hydroxy-3-methoxyphenyl)-1,6-heptadiene-3,5-dione] in the rat. *Xenobiotica*, *8*: 761–768, 1978.
37. Ravindranath, V., and Chandrasekhara, N. Absorption and tissue distribution of curcumin in rats. *Toxicology*, *16*: 259–265, 1980.
38. Shoba, G., Joy, D., Joseph, T., Majeed, M., Rajendran, R., and Srinivas, P. S. S. R. Influence of piperine on the pharmacokinetics of curcumin in animals and human volunteers. *Planta Med.*, *64*: 353–356, 1998.
39. Pan, M. H., Huang, T. M., and Lin, J. K. Biotransformation of curcumin through reduction and glucuronidation in mice. *Drug Metab. Dispos.*, *27*: 486–494, 1999.
40. Pool-Zobel, B. L., Abrahamse, S. L., Collins, A. R., Kark, W., Gugler, R., Oberreuther, D., Siegel, E. G., Treptow van Lishaut, S., and Rechkemmer, G. Analysis of DNA strand breaks, oxidized bases, and glutathione *S*-transferase P1 in human colon cells from biopsies. *Cancer Epidemiol. Biomark. Prev.*, *8*: 609–614, 1999.
41. Oetari, S., Sudibyo, M., Commandeur, J. N. M., Samhøedi, R., and Vermeulen, N. P. E. Effect of curcumin on cytochrome P450 and glutathione *S*-transferase activities in rat liver. *Biochem. Pharmacol.*, *51*: 39–45, 1996.
42. Piper, T., Singhal, S. S., Salameh M., Torman, R. T., Awasthi, Y. C., and Awasthi, S. Mechanisms of anticarcinogenic properties of curcumin: the effect of curcumin on glutathione linked detoxification enzymes in rat liver. *Int. J. Biochem. Cell Biol.*, *30*: 445–456, 1998.
43. Susan, M., and Rao, M. N. A. Induction of glutathione *S*-transferase activity by curcumin in mice. *Drug Res.*, *42*: 962–964, 1992.
44. Nijhoff, W. A., Groen, G. M., and Peters, W. H. M. Induction of rat hepatic and intestinal glutathione *S*-transferases and glutathione by dietary naturally occurring anticarcinogens. *Int. J. Oncol.*, *3*: 1131–1139, 1993.

Clinical Cancer Research

Effects of Dietary Curcumin on Glutathione S-Transferase and Malondialdehyde-DNA Adducts in Rat Liver and Colon Mucosa: Relationship with Drug Levels¹

Ricky A. Sharma, Christopher R. Ireson, Richard D. Verschoyle, et al.

Clin Cancer Res 2001;7:1452-1458.

Updated version Access the most recent version of this article at:
<http://clincancerres.aacrjournals.org/content/7/5/1452>

Cited articles This article cites 39 articles, 21 of which you can access for free at:
<http://clincancerres.aacrjournals.org/content/7/5/1452.full.html#ref-list-1>

Citing articles This article has been cited by 20 HighWire-hosted articles. Access the articles at:
</content/7/5/1452.full.html#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, contact the AACR Publications Department at permissions@aacr.org.